

Wavelet-Based Functional Mixed Models

Jeffrey S. Morris

**UT MD Anderson Cancer Center
Houston, Texas**





Functional Data

- **Functional Data:**
 - Ideal units of observation: **curves**
 - Observed data: **curves sampled on fine grid**
- Increasingly encountered in biomedical research with new technologies taking automated measurements
- Present unique challenges:
 - Extremely **large data sets** (>100s-1000s per curve)
 - Curves may be **complex** and **irregular**, spatially heterogeneous with many local features

Example: Accelerometer Data

- **Accelerometers:** small motion sensors that digitally record minute-by-minute activity levels
 - Increasingly used in surveillance and intervention studies
- **TriTrac-R3D:** sensor worn on hip
 - Minute-by-minute record of motion in 3 planes
 - Condensed into single activity level measurement/minute
 - Activity “profile” for each day

Accelerometer Data



Accelerometer Data

- **Planet Health Study** (Harvard University):
 - Intervention study investigating activity levels of middle school children in Boston area schools
 - Children's activity levels objectively monitored using TriTrac-R3D activity monitor for one or two 4-day sessions
 - **Data considered:** 292 daily profiles/103 children/5 schools
660 measurements/profile (every minute from 9am-8pm)
- **Goals:**
 1. Assess how activity levels vary throughout day, across schools, across different days of the week, over time from early to late Spring, and across various child-level covariates.
 2. Assess relative variability in activity levels from day-to-day and child-to-child, in order to guide future study design.

Linear Mixed Models

Linear Mixed Model (Laird and Ware, 1982):

$$\underbrace{Y}_{N \times 1} = \underbrace{X}_{N \times p} \underbrace{\beta}_{p \times 1} + \underbrace{Z}_{N \times m} \underbrace{u}_{m \times 1} + \underbrace{e}_{N \times 1}$$

$$\begin{aligned} u &\sim N(0, \overbrace{D}^{m \times m}) \\ e &\sim N(0, \underbrace{R}_{N \times N}) \end{aligned}$$

- **Fixed effects** part, $X\beta$, accommodate a broad class of mean structures, including main effects, interactions, and linear coefficients.
- **Random effects** part, Zu , provide a convenient mechanism for modeling correlation among the N observations.

Functional Mixed Model

Suppose we observe a sample of N curves,
 $Y_i(t)$, $i=1, \dots, N$, all defined on \mathcal{T}

$$U_k(t) \sim GP(0, Q)$$

$$E_i(t) \sim GP(0, S)$$

$$Y_i(t) = \sum_{j=1}^p X_{ij} B_j(t) + \sum_{k=1}^m Z_{ik} U_k(t) + E_i(t)$$

- $B_j(t)$ = fixed effect functions
- $U_k(t)$ = random effect functions
- $E_i(t)$ = residual error processes
- Q and S are covariance surfaces on $\mathcal{T} \times \mathcal{T}$
 - $S(t_1, t_2) = \text{Cov}\{E_i(t_1), E_i(t_2)\}$: describes **within-curve** covariance structure of residual curve-to-curve deviations

Functional Mixed Models

- **Key feature of FMM:** Does not require specification of parametric form for curves
- **Kernels/fixed-knot splines** may not work well for spatially heterogeneous data – inherent smoothness assumptions attenuate local features
- **Wavelet Regression:** nonparametric regression technique that better preserves local features present in the curves.

Introduction to Wavelets


- **Wavelets:** families of orthonormal basis functions

$$g(t) = \sum_{j,k \in \mathfrak{I}} d_{jk} \psi_{jk}(t)$$

$$\psi_{jk}(t) = 2^{-j/2} \psi(2^{-j/2} t - k)$$

$$d_{jk} = \int g(t) \psi_{jk}(t) dt$$

Daubechies (4) Basis Function



- **Discrete Wavelet Transform (DWT):** fast algorithm $\{O(T)\}$ for obtaining T empirical wavelet coefficients for curves sampled on equally-spaced grid of length T .
- **Linear Representation:** $d = y W'$
 - W' = T -by- T orthogonal projection matrix
- **Inverse DWT (IDWT):** $y = d W$

Wavelet Regression

- **Wavelet Regression** – 3 step process
 1. Project data into wavelet space
 2. Threshold/shrink coefficients
 3. Project back to data space
- Yields *adaptively regularized* (plot) nonparametric estimates of function
- Morris, et al. (2003) extended to hierarchical functional model (Bayesian)
- Morris and Carroll (2006) extended to general functional mixed model framework (wavelet-based functional mixed model)

Functional Mixed Model

(Discrete version)

Y = **N-by-T matrix** containing the **observed spectra** on sampling grid of size **T**

$$\underbrace{Y}_{N \times T} = \underbrace{X}_{N \times p} \underbrace{B}_{p \times T} + \underbrace{Z}_{N \times m} \underbrace{U}_{m \times T} + \underbrace{E}_{N \times T}$$
$$U_i \sim MVN(0, Q)$$
$$E_i \sim MVN(0, S)$$

- B_{ij} is the effect of covariate i at location t_j
- Q and S are covariance matrices ($T \times T$)
- Note: Some structure must be assumed on form of Q and S (discussed later)

Wavelet-Based FMM: General Approach

1. **Project** observed functions Y **into wavelet space.**
2. **Fit FMM** in wavelet space.
(Use MCMC to get posterior samples)
3. **Project** wavelet-space estimates
(posterior samples) **back to data space.**

Wavelet-Based FMM: General Approach

- 1. Project** observed functions **Y** **into** wavelet space.
- 2. Fit FMM** in wavelet space
(Use MCMC to get posterior samples)
- 3. Project** wavelet-space estimates
(posterior samples) **back to data space.**

Wavelet-Based FMM

1. Project observed functions Y to wavelet space

- Apply DWT to rows of Y to get wavelet coefficients corresponding to each observed function

$$\underbrace{D}_{N \times T} = \underbrace{Y}_{N \times T} \underbrace{W'}_{T \times T}$$

- Projects the observed curves into the space spanned by the wavelet bases.

Wavelet-Based FMM:

General Approach

1. **Project** observed functions **Y** **into wavelet space.**
2. **Fit FMM** in wavelet space
(Use MCMC to get posterior samples)
3. **Project** wavelet-space estimates
(posterior samples) **back to data space.**

Projecting FMM to Wavelet Space

$$\underbrace{Y}_{N \times T} = \underbrace{X}_{N \times p} \underbrace{B}_{p \times T} + \underbrace{Z}_{N \times m} \underbrace{U}_{m \times T} + \underbrace{E}_{N \times T}$$

$$U_i \sim MVN(0, Q)$$

$$E_i \sim MVN(0, S)$$

Projecting FMM to Wavelet Space

$$\underbrace{Y}_{N \times T} \underbrace{W'}_{T \times T} = \underbrace{X}_{N \times p} \underbrace{B}_{p \times T} + \underbrace{Z}_{N \times m} \underbrace{U}_{m \times T} + \underbrace{E}_{N \times T}$$

$$U_i \sim MVN(0, Q)$$

$$E_i \sim MVN(0, S)$$

Projecting FMM to Wavelet Space

$$\underbrace{Y}_{N \times T} \underbrace{\mathbf{W}'}_{T \times T} = \underbrace{X}_{N \times p} \underbrace{B}_{p \times T} \underbrace{\mathbf{W}'}_{T \times T} + \underbrace{Z}_{N \times m} \underbrace{U}_{m \times T} \underbrace{\mathbf{W}'}_{T \times T} + \underbrace{E}_{N \times T} \underbrace{\mathbf{W}'}_{T \times T}$$

$$U_i \sim MVN(0, Q)$$

$$E_i \sim MVN(0, S)$$

Projecting FMM to Wavelet Space

$$\underbrace{Y}_{N \times T} \underbrace{W'}_{T \times T} = \underbrace{X}_{N \times p} \underbrace{B}_{p \times T} \underbrace{W'}_{T \times T} + \underbrace{Z}_{N \times m} \underbrace{U}_{m \times T} \underbrace{W'}_{T \times T} + \underbrace{E}_{N \times T} \underbrace{W'}_{T \times T}$$

$$U_i W' \sim MVN(0, W Q W')$$

$$E_i W' \sim MVN(0, W S W')$$

Projecting FMM to Wavelet Space

$$\underbrace{\mathbf{D}}_{N \times T} = \underbrace{\mathbf{X}}_{N \times p} \underbrace{\mathbf{B}^*}_{p \times T} + \underbrace{\mathbf{Z}}_{N \times m} \underbrace{\mathbf{U}^*}_{m \times T} + \underbrace{\mathbf{E}^*}_{N \times T}$$

$$\mathbf{U}_i^* \sim MVN(0, \mathbf{Q}^*)$$

$$\mathbf{E}_i^* \sim MVN(0, \mathbf{S}^*)$$

Prior Assumptions

Mixture prior on β_{ijk}^* :

$$\beta_{ijk}^* = \gamma_{ijk}^* N(0, \tau_{ij}) + (1 - \gamma_{ijk}^*) \delta_0$$

$$\gamma_{ijk}^* = \text{Bernoulli}(\pi_{ij})$$

- Nonlinearly shrinks β_{ijk}^* towards 0, leading to **adaptively regularized** estimates of $\beta_i(t)$.
- τ_{ij} & π_{ij} are **regularization parameters**
 - Can be estimated from the data using **empirical Bayes**
 - Extend Clyde&George (1999) to functional mixed model

Model Fitting

- **MCMC** to obtain posterior samples of model quantities
 - Work with marginal likelihood; U^* integrated out;
- Let Ω be a vector containing ALL covariance parameters (i.e. Q^* and S^*).

MCMC Steps

1. Sample from $f(B^*/D, \Omega)$:

Mixture of normals and point masses at 0 for each i, j, k .

2. Sample from $f(\Omega/D, B^*)$:

Metropolis-Hastings steps for each j, k

3. If desired, sample from $f(U^*/D, B^*, \Omega)$:

Multivariate normals

Wavelet-Based FMM: General Approach

1. **Project** observed functions Y **into wavelet space.**
2. **Fit FMM** in wavelet space
(Use MCMC to get posterior samples)
3. **Project** wavelet-space estimates
(posterior samples) **back to data space.**

Wavelet-Based FMM

3. **Project** wavelet-space estimates (posterior samples) **back to data space**.

- Apply IDWT to posterior samples of B^* to get posterior samples of fixed effect functions $B_j(t)$ for $i=1, \dots, p$, on grid t .
 - **$B=B^*W$**
- Posterior samples of $U_k(t)$, Q , and S are also available, if desired.
- Can be used for Bayesian inference/prediction

Model


Let Y be 292×660 matrix containing 292 accelerometer profiles for each minute from 9am-8pm.

$$Y = XB + ZU + E$$

- $X = 292 \times 14$ matrix of covariates
 - School effects (5), gender, triceps calipers, BMI, day-of-week (4), daylight savings time, avg tv hrs/wk
- $B = 14 \times 660$ matrix of fixed effects functions
 - B_{ij} is effect of covariate i at time t_j
- $Z = 292 \times 106$ matrix indicating child for each profile
 $U = 106 \times 660$ matrix of random effect functions (1/child)
- $E = 292 \times 660$ matrix of residual errors

Selected Results: **School Effects**

(a) School E




- **School schedules evident in effects**
 - Spikes every 48min (changing classes)
 - 3 lunch periods
 - School out at 2:15pm
- **Not so evident in individual curves**

4/4/2007


<http://biostatistics.mdanderson.org/Morris> (20min)

Selected Results: DST Effect



- DST – April 6th
- **More active after DST** (overall 8%, $p=0.062$)
- Especially strong:
 - **As school is letting out** (2:15-3:00, 25%, $p=0.03$)
 - **In early evening** (5:30-7:00, 30%, $p=0.01$)
- Note: Sunset was
 - 5:10-6:15 before DST
 - 7:15-8:10 after DST

Selected Results: **TV hours/wk**



- **TVhrs coded as continuous factor (standardized)**
- **TVhrs effect negative (-1.3% per sd, $p=0.03$)**
 - More TV, less active
 - 3:00-5:30, -2.6%, $p=0.02$
 - 7:00-8:00, -3.6%, $p=0.008$
- **Positive effect over lunch**
 - +2.7%, $p=0.03$
 - More TV, on average more active over lunch

Results: Covariance Analysis

- **Variability: 91% day-to-day, 9% child-to-child**
 - Important to have many days per child
- **Study variability as function of t**
 - **Child-to-child** variability: **school day > after school**
 - **Day-to-day** variability: **after school > school day**
 - Relative day-to-day variability after school: **95%-99%**
- **Equivalent designs:**
 - **108** children, **4** days/child
 - **72** children, **8** days/child
 - **54** children, **16** days/child
- **Less children, more days, save \$\$\$?**

Discussion

- Introduced unified modeling approach for FDA
 - Can accommodate very irregular functions
- Method based on mixed models; is **FLEXIBLE**
 - Accommodates a **wide range of experimental designs**
 - Addresses **large number of research questions**
- Posterior samples allow **Bayesian inference and prediction**
 - **Posterior credible intervals**; pointwise or joint
 - **Predictive distributions** for future sampled curves
 - **Predictive probabilities** for classification of new curves
 - Bayesian functional inference can be done via **Bayes Factors**
- Since a unified modeling approach is used, all **sources of variability** in the model **propagated throughout inference**.

Discussion

- Approach is Bayesian. The **only informative priors to elicit are regularization parameters**, which can be estimated from data using empirical Bayes.
- Developed **general-use code** – reasonably fast and straightforward to use → minimum information to specify is Y, X, Z matrices.
- Can deal with **missing data**, i.e. partially observed functions (not discussed here)
- Method **generalizes to higher dimensional functions**, e.g. image data, space/time (fixed domain) data.

Acknowledgements

- Work presented here is from 2 papers
 1. “*Wavelet-Based Functional Mixed Models*” (2006) Jeffrey S. Morris and Raymond J. Carroll, *JRSS-B*, 68(2): 179-199.
 2. “*Using Wavelet-Based Functional Mixed Models to Characterize Population Heterogeneity in Accelerometer Profiles: A Case Study*” (2006) Jeffrey S. Morris, Cassandra Arroyo, Brent Coull, Louise Ryan, Richard Herrick, and Steve Gortmaker, *JASA*, to appear.
 3. “*Bayesian Analysis of Mass Spectrometry Proteomics Data using Wavelet Based Functional Mixed Models*” (2006) Jeffrey S. Morris, Philip J. Brown, Richard Herrick, Keith A. Baggerly, and Kevin R. Coombes, submitted, available on MDACC’s bepress site.
- Computer code/papers on web at <http://biostatistics.mdanderson.org/Morris/papers.html>


Accelerometer Example

Let $Y_{ij}(t)$ be accelerometer profile on day j from child i

$$Y_{ij}(t) = B_0(t) + \sum_{k=1}^{p_1} X_{ik} B_k^{child}(t) + \sum_{k=1}^{p_2} X_{ijk} B_k^{day}(t) + U_i(t) + E_{ij}(t)$$


- X_{ik} = child-level covariates (school, race, gender, BMI, % body fat, avg hrs of TV/week)
- X_{ijk} = day-level covariates (day-of-week, DST)
- $B_0(t)$ = overall mean profile
- $B_k^{child}(t)$ = effect functions for child-level covariates
- $B_k^{day}(t)$ = effect functions for day-level covariates
- $U_i(t)$ = Random effect function for child i

Results: Bouts







- Can compute posterior predictive probabilities of bouts for children

Results: Bouts



- Can compute posterior predictive probabilities of bouts for children
- **Model-based** predictive probabilities not far from **empirically-estimated** probabilities
- May want heavier tails

Some Results



- **School #5:**

- Spikes every 48min (changing classes)
- 3 lunch periods
- School out at 2:15pm

- **BMI effect positive**

- Artifact of preprocessing?


- **Daylight Savings Time**

- More active after DST
- Especially 2-3pm, 5-7pm

- **Avg hrs TV/week**

- More TV=less active
- Especially 3-5pm, 7-8pm
- More active at lunch

Results




4/4/2007


<http://biostatistics.mdanderson.org/Morris>

Adaptive Regularization


Regularization by Local Linear Smoothing



Adaptive Regularization by Wavelet Shrinkage




Simulation: Covariance Structure



- **True mean:** line plus peak
- **True variance:** increasing in t , with extra var at peak
- **True autocorrelation:** Strong autocorrelation (0.9) at left, weak autocorrelation (0.1) right, extra at peak

Simulation: Covariance Structure



- **Independence in wavelet space** accommodates varying degrees of **autocorrelation in data space**
- Allowing variance components to vary across scale j and location k accommodates **nonstationarities**


Model Each Column Separately

$$\underbrace{d_{jk}}_{N \times 1} = \underbrace{X}_{N \times p} \underbrace{\beta_{jk}^*}_{p \times 1} + \underbrace{Z}_{N \times m} \underbrace{u_{jk}^*}_{m \times 1} + \underbrace{e_{jk}^*}_{N \times 1}$$

$$u_{jk}^* \sim N(\mathbf{0}, q_{jk}^*)$$

$$e_{jk}^* \sim N(\mathbf{0}, s_{jk}^*)$$

Selected Results: **BMI Effect**



- BMI Coded as continuous factor (mean-centered)
- BMI effect positive ($p < 0.0005$)
 - Higher BMI, more active
 - Preprocessing artifact?
- Should raw activity levels be monitored instead of METs?